

Technical Note

Establishing a cell-free *Vibrio natriegens* expression system

Daniel Wiegand, Henry H Lee, Nili Ostrov, and George M. Church

ACS Synth. Biol., **Just Accepted Manuscript** • DOI: 10.1021/acssynbio.8b00222 • Publication Date (Web): 30 Aug 2018Downloaded from <http://pubs.acs.org> on September 1, 2018

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



Establishing a cell-free *Vibrio natriegens* expression system

Authors: Daniel J. Wiegand^{1,2}, Henry H. Lee¹, Nili Ostrov^{1*}, George M. Church^{1,2*}

Author affiliation:

¹ Department of Genetics, Harvard Medical School, Boston, MA 02115

² Wyss Institute for Biologically Inspired Engineering, Boston, MA 02115

* Correspondence to: nili_ostrov@hms.harvard.edu, gchurch@genetics.med.harvard.edu

Abstract

The fast growing bacterium *Vibrio natriegens* is an emerging microbial host for biotechnology. Harnessing its productive cellular components may offer a compelling platform for rapid protein production and prototyping of metabolic pathways or genetic circuits. Here, we report the development of a *V. natriegens* cell-free expression system. We devised a simplified crude extract preparation protocol and achieved >260 µg/mL of super-folder GFP in a small-scale batch reaction after three hours. Culturing conditions, including growth media and cell density, significantly affect translation kinetics and protein yield of extracts. We observed maximal protein yield at incubation temperatures of 26°C or 30°C, and show improved yield by tuning ions crucial for ribosomal stability. This work establishes an initial *V. natriegens* cell-free expression system, enables probing of *V. natriegens* biology, and will serve as a platform to accelerate metabolic engineering and synthetic biology applications.

Keywords: *Vibrio natriegens*, crude cell extract, cell-free system, TX-TL, *in-vitro* protein synthesis

1
2
3 With the shortest doubling time of all known organisms, *Vibrio natriegens* has garnered
4 considerable interest as a promising microbial host to accelerate research and biotechnology¹⁻⁴.
5 Its rapid growth rate, which has been linked to high rates of protein synthesis⁵ and metabolic
6 efficiency^{6,7} suggests that this host may be harnessed as a powerful cell-free expression system.
7 Cell-free bioproduction of protein and chemicals has been extensively investigated in *E. coli* and
8 recently expanded to several other bacteria⁸⁻¹³. Accordingly, we assessed *V. natriegens* crude
9 cell extract productivity in small-scale batch reactions using super-folder GFP (sfGFP)
10 controlled by a T7 promoter. We systematically explored culturing conditions and extract
11 additives to provide a simple and robust *V. natriegens* cell-free system. Our calibration offers
12 initial conditions for cell-free protein expression with *V. natriegens* and sheds light on critical
13 factors for future process engineering^{14,15}
14
15
16
17
18

19 To begin developing a cell-free protocol for *V. natriegens*, we evaluated established methods for
20 extract preparation for *E. coli* and other bacterial systems^{10,16-18}. Overall, we aimed for general
21 accessibility and reproducibility by opting at each step for commonly available equipment while
22 considering ease of user operation and cost reduction. Our preparation protocol utilizes one liter
23 cultures in shake flasks, cell lysis by pulse sonication, and small-scale (10 μ L) batch reactions in
24 a 96- or 384-well format to maximize parallelization and screening throughput¹⁷.
25
26
27

28 We first assessed the effect of *V. natriegens* culturing conditions on extract productivity. We
29 tested commonly available microbiological media, supplemented with salt as needed to maintain
30 robust growth^{1,2,5,19}. Crude cell extracts were prepared from *V. natriegens* cultures growing in
31 the following six media: LB with 3% (w/v) NaCl (LB3), LB with V2 salts (LB-V2), LB with
32 Ocean Salts (LBO), Nutrient Broth with Ocean Salts (NBO), Brain Heart Infusion Broth with
33 Ocean Salts (BHIO), and Marine Broth (MB). Cultures were grown at 30°C and harvested at
34 OD₆₀₀ = 1.0 (**Methods**).
35
36
37
38

39 We found that the choice of growth media significantly affects cell extract productivity. Extracts
40 cultured in LB-V2 produced the highest protein yield (196 \pm 12.46 μ g/mL), followed by extracts
41 cultured in BHIO (58.86 \pm 2.61 μ g/mL) and LB3 (33.18 \pm 5.57 μ g/mL) (**Figure 1a**,
42 **Supplementary Figure 1a**). Low protein yield was observed from extracts cultured in NBO and
43 LBO media, with no significant protein expression observed from extracts cultured in MB
44 media. We thus elected to use LB-V2 extracts for all further investigations.
45
46
47

48 Previous studies have suggested that higher cellular ribosomal content is required to support
49 faster growth rates^{5,20}. To test whether extract productivity could be used for a proxy for active
50 ribosomes in each growth medium, we assessed the correlation between the culture generation
51 time and extract protein yield (**Figure 1b**, **Supplementary Figure 1c**). Our data shows no
52 relationship between generation time and cell extract productivity. The three shortest generation
53 times at 30°C were MB, LB-V2, and LB3 (20.14 \pm 2.16, 20.55 \pm 0.63 and 22.14 \pm 1.13
54
55
56
57
58
59
60

1
2
3 minutes, respectively). However, extracts from LB-V2 cultures produced significantly more
4 protein than extract from all other culture media (~6-fold more than LB3).
5
6

7 To gain deeper insight into the difference between growth media, we assessed translation
8 kinetics for each growth media type by examining the rate of accumulating sfGFP (**Figure S1b,**
9 **Supplementary Figure 1b, Methods**). As expected, LB-V2 extracts had the maximum rate of
10 protein synthesis, which was achieved after ~30 minutes. However, NBO extracts reached its
11 maximum synthesis rate in only ~15 minutes, although the rate was 2.6-fold lower, and the total
12 protein yield was 19.6-fold lower. Intriguingly, although their rate and total yield were
13 significantly lower than that of LB-V2 extract, BHIO and LB3 extracts appear to sustain protein
14 synthesis more robustly compared to other extracts, as evidenced by the protracted time
15 constants for maximum and minimal rates. All extracts demonstrated a decay in rate of protein
16 expression after 60 minutes, indicating consumption of input building blocks, buildup of
17 inhibitory byproducts, or depletion of the sfGFP template. Overall, these results indicate
18 significant variation in the fraction of active translation between media types. Specifically,
19 potassium and magnesium ions, which are present in LB-V2 but not LB3, may enhance the
20 stability of translation components in crude cell extract. Given the significant impact of culture
21 media on extract yield, further investigation and development of customized cell-free growth
22 medium for extract preparation may be required.
23
24
25
26
27
28

29 Additionally, we tested whether a post-lysis run-off reaction, previously shown to increase the
30 availability of translational machinery in crude cell extracts¹⁷, would be beneficial to increasing
31 protein yield. We performed run-off reactions prior to addition of the sfGFP template, as
32 previously reported^{21,22}. We found that run-off reactions resulted in a significantly reduced
33 capacity to produce sfGFP (**Supplementary Figure 8**).
34
35
36

37 Active cell-free extracts are routinely produced using exponentially growing cells^{17,23}. We thus
38 tested the impact of *V. natriegens* cell density at harvest on extract productivity. Cells were
39 grown in LB-V2 media, harvested at OD₆₀₀ of 0.5, 1.0 or 3.0 and the corresponding sfGFP yield
40 was measured (**Figure 1c, Supplementary Figure 2**). We found the most productive extracts
41 were obtained from mid-logarithmic cultures (OD₆₀₀ = 1.0), compared to 6.2-fold lower yield
42 from late-logarithmic cultures (OD₆₀₀ = 3.0). We were unable to detect sfGFP production from
43 extracts obtained from early-logarithmic cultures (OD₆₀₀ = 0.5). These effects of cell density on
44 extract performance are consistent with those reported for other cell-free systems¹⁷.
45
46
47
48

49 Cell-free extracts uniquely allow cellular components to operate decoupled from their culturing
50 conditions. We next turned to calibrating cell-free reaction conditions. We examined incubation
51 temperatures ranging 18-37°C and observed maximal protein expression at 26°C and 30°C
52 (262.28 ± 8.13 µg/mL and 265.01 ± 8.15 µg/mL, respectively) (**Figure 2a**). While *in vivo* growth
53 rate for *V. natriegens* has been found to increase with temperature and to be maximal at 37°C
54
55
56
57
58
59
60

1
2
3^{1,19}, we found that extracts incubated at 30°C produced 3-fold more protein than 37°C. These
4 results support the hypothesis that the *V. natriegens*' rapid growth at a higher temperature is
5 aided by a larger number of ribosomes, rather than more efficient ones⁵. It may also be the case
6 that more functional ribosomes are present at lower temperatures despite a higher total number at
7 higher temperature as previously observed in *E. coli*²⁴. Further investigation of ribosome content
8 and availability is warranted to elucidate underlying biology. In addition, the reduced protein
9 yields at temperatures higher than 30°C may be artifactual due to our choice of T7 polymerase.
10 Due to its high processivity, T7 polymerase has been shown to reduce protein yield for a
11 bacterial cell-free system by disrupting the coupling of transcription and translation²⁵. Further
12 improvements may be made by using mutant T7 polymerases or endogenous promoters. For ease
13 of further investigations, we chose to incubate all *V. natriegens* cell-free reactions at 26°C.
14
15
16
17
18

19 Having set an incubation temperature for optimal extract productivity, we then tuned the range of
20 K⁺ and Mg²⁺ ions in the extract reaction^{10,18}. In general, we found the addition of K⁺ had a
21 significant effect on both yield and rate of protein expression, while Mg²⁺ had only moderate
22 effect at the tested conditions (**Figure 2b-c, Supplementary Figure 3**). We set our assay
23 conditions at 80 mM K⁺ and 3.5 mM Mg²⁺, which yielded the highest reaction productivity for
24 sfGFP expression (248.75 ± 6.82 μg/mL and 217.87 ± 23.40 μg/mL, respectively). Notably,
25 while higher ion concentrations have been reported²⁶, the determined optimal ion concentrations
26 for both K⁺ and Mg²⁺ in our system are consistent with other cell-free systems which have been
27 developed with 3-Phosphoglyceric acid (3-PGA) as an energy source^{10,18,27}.
28
29
30
31

32 We then sought to determine if the total amount of crude extract needed per sample could be
33 lowered while retaining reaction productivity. The percent of cell-free reaction that comprises
34 crude cell extract varies between different bacterial systems^{12,17} and titration of the cellular
35 components has been shown to improve protein yield⁹. We thus tested reactions where the
36 percent of extract was varying, ranging 35% to 15%, and monitored the kinetics of sfGFP
37 accumulation. Surprisingly, we found that decreasing the extract from 35% to 30% of the total
38 reaction volume resulted in significant improvement in protein yield (43% increase) with no
39 significant change in expression rate (**Figure 2d, Supplementary Figure 4**). Further decrease in
40 extract amount (from 30% to 25%) resulted in a decreased rate of protein expression without
41 significant change in protein yield. These results indicate up to half of the original extract
42 volume can be used with increased or equal reaction productivity, which increases the number of
43 cell-free reactions produced from a single culture batch. It is worth also noting that the lowering
44 the total extract percentage per reaction volume yielded sustained protein production for the
45 duration of the reaction (**Supplementary Figure 4**).
46
47
48
49
50
51

52 The use of linear, rather than plasmid, template DNA in cell-free expression systems is desirable
53 for high-throughput and rapid testing¹⁸. We thus evaluated protein production from a linear PCR
54 template compared to a circular plasmid template (**Figure 2e**). In addition to sfGFP under T7
55
56
57
58
59
60

1
2
3 promoter, our PCR amplicon also contained 58bp of non-coding DNA at the 3' end of the
4 construct. We observed protein production from the PCR amplicon to be 13.5-fold lower than an
5 equimolar amount of circular plasmid, suggesting rapid DNA degradation by cellular nucleases.
6 To test whether a linear template could be protected from nuclease digestion, we assessed protein
7 yield from a PCR product with two phosphorothioated bonds on each end²⁴. Notably, this
8 modified template was also 58bp shorter, carrying less non-coding sequence at the 3'. We
9 observed no improvement in protein yield using protected PCR amplicon (**Figure 2e**), suggestive
10 of endonuclease, rather than exonuclease, activity. The modified short template had 2-fold lower
11 yield compared with longer, non-modified template. Padding of linear templates with non-coding
12 sequences²⁸ or inactivation of endogenous nucleases by genome engineering will likely improve
13 protein yield significantly and warrants further investigation.
14
15
16
17
18

19 To monitor the kinetics of linear DNA template degradation, we used a fluorescence-based assay
20 to quantify the amount of intact linear DNA template in the cell-free reaction over time
21 (**Methods**)²⁸. We found that more than 50% of linear DNA template was degraded by 10
22 minutes of co-incubation with the cell-free extract (**Figure 2f**). No significant difference in
23 degradation was observed using the shorter phosphorothioated template, corroborating our
24 previous observations that protecting the 5' ends of linear DNA does not improve template
25 stability and that less non-coding padding decreases protein yield.
26
27
28
29

30 Previously, it was shown that linear template stability can be increased in *E.coli* cell-free system
31 with inclusion of the lambda phage protein GamS, which inhibits the host RecBCD nuclease
32 complex^{28,29}. However, we observed no change in the extent of linear template degradation upon
33 addition of lambda GamS protein to *V. natriegens* cell-free reactions (**Supplementary Figure 9**).
34 The same results were observed for both phosphorothioate protected and unprotected linear
35 templates. This could indicate that GamS does not interact with the DNA-binding domain of *V.*
36 *natriegens*' RecBCD exonuclease complex or that linear template degradation occurs by an
37 independent mechanism. Alternatively, degradation may result from endonuclease activity given
38 that the inclusion phosphorothioate bonds at both the 5' and 3' termini the linear template did not
39 alter degradation kinetics (**Figure 2f**). Additional research is required to elucidate *V. natriegens*
40 exo- or endonuclease and effective strategies to inhibit their activity. Engineering of the host
41 genome to remove nucleases may enable facile production of highly active extracts for high
42 efficiency protein expression with linear templates.³⁰
43
44
45
46
47

48 Maximizing protein expression in cell-free systems requires the combinatorial optimization of a
49 large parameter search space and no single best approach currently exists^{21,22,31,32}. During this
50 work, an alternative *V. natriegens* CFPS has been reported²⁶, which explores the use of other
51 culture medium, energy regeneration sources, and extract preparation. For example, while our
52 screen of growth media shows that the use of LB-V2, rather than BHIO, results in higher protein
53 expression yields (**Supplementary Figure 1**), the alternative report showed that an extensive
54
55
56
57
58
59
60

1
2
3 purification process can provide more cell-free protein expression in cells harvested from BHIO
4 culture media at high optical density (OD=2.5). Additionally, we found that run-off did not
5 promote increased protein expression, which may indicate altered ribosomal occupancy under
6 different culture conditions or growth stage at the time of harvest. Consistent with previous
7 reports^{33–35}, it is interesting to note the higher concentration of Mg²⁺ required when using
8 creatine phosphate (CP) as energy regeneration system compared with 3-PGA in this work.
9 Additional investigation will be essential to fully develop the potential of *V. natriegens* for
10 CFPS. While protein yield is an important feature, it may not be the sole consideration when
11 working with CFPS. The protocol for extract preparation presented in this work does not require
12 specialized equipment or lengthy purification steps, and is therefore short, accessible, and simple
13 to execute, which should facilitate further experimentation.
14
15
16
17
18

19 Throughout the development of this preparation protocol for *V. natriegens* CFPS, we have
20 utilized the same methods and reagents, where appropriate, to prepare extracts from *E. coli* A19,
21 an RNase I- deficient strain that is commonly used for cell-free reactions, to validate protein
22 expression functionality^{36,37}. Crude cell extract was prepared as previously described from *E.*
23 *coli* cultured in rich media (YPG) at 37°C²³ and harvested at OD₆₀₀=1.0 (**Methods**). We then
24 performed similar calibration of reaction conditions for *E.coli* as described above for *V.*
25 *natriegens* (**Supplementary Figure 5**). For context, we observed comparable protein yields
26 between *E.coli* A19 and wild-type *V. natriegens* (**Supplementary Figure 6**) when production
27 occurred in their respective optimal reaction conditions. The yield obtained from *E. coli* A19
28 extracts are similar to previously reported GFP yields using small-batch conditions³⁸, which
29 supports the robustness of our protocol. Kinetic analyses indicated that *V. natriegens* extracts
30 sustained elevated protein expression rates over 60 minutes, while *E. coli* extracts only sustain
31 elevated rates for 20 minutes. More broadly, this system produces more protein than cell-free
32 systems derived from *Bacillus subtilis*²⁷ and *Streptomyces venezuelae*¹⁰, as measured by sfGFP
33 yield (~9.3 μM sfGFP in our 10 μL *V. natriegens* cell-free reaction, compared with 0.8 μM and
34 1.3 μM, respectively).
35
36
37
38
39
40

41 Taken together, these results provide a foundation for cell-free protein expression with wild-type
42 *V. natriegens* and may direct further efforts to establish designated cell-free strains. Higher
43 protein yield could be achieved by further optimization of culturing and preparation conditions
44 and may include: semi-continuous extract reactions to allow for energy regeneration, resupplying
45 of amino acids, and the removal of waste products^{12,39}. In addition, engineering of *Vibrio*
46 *natriegens*, for example using DNase- or RNase-deficient strains, removal of deleterious and
47 competing metabolic pathways, and expression of additional tRNAs will likely further enhance
48 protein yield from extracts⁴⁰. These additional improvements will also enable the use of linear
49 DNA to facilitate rapid prototyping and high-throughput screens. This work establishes an initial
50 cell-free system in *V. natriegens* and sheds light on important aspects for continued
51 development.
52
53
54
55
56
57
58
59
60




Figure 1. *V. natriegens* culture conditions impact extract protein yield. (a) Effect of culture growth media on cell-free extract protein yield. *V. natriegens* cells were grown in the indicated media at 30°C to OD₆₀₀ = 1.0. Crude cell lysate was prepared production of sfGFP measured over three hours. (LB3 - LB with 3% (w/v) NaCl; LB-V2 - LB with V2 salts; LBO - LB with Ocean Salts, NBO - Nutrient Broth with Ocean Salts; BHIO - Brain Heart Infusion with Ocean Salts; and MB - Marine Broth). (b) Generation time and sfGFP yield for each media type tested. (c) Optical density at the time of cell harvest. Unless otherwise indicated, cell-free reactions were incubated at 26°C in a thermocycler using 500 ng of plasmid DNA, 160 mM K-glutamate and 3.5 mM Mg-glutamate. sfGFP yield was measured after 3 hours. The mean and standard deviations are shown (N=3).




Figure 2. Properties of *V. natriegens* crude cell-free extract. (a) Cell-free incubation temperature. (b) Supplemented potassium ions in reaction buffer. (c) Supplemented magnesium ions in reaction buffer. (d) Percent of extract used relative to total reaction volume. (e) Template DNA provided in the cell-free reaction. Equimolar amount of circular plasmid (pJL1, grey), linear DNA (PCR product, black solid line) or linear DNA with two phosphorothioated bonds on each end (PCR product, black dotted line) was used as template for cell-free expression of sfGFP. (f) Degradation of linear DNA template. Fluorescence of AlexaFluor 594-5-dUTP labeled DNA template was monitored over two hours. Linear template with (dotted) or without (solid) two phosphorothioated bonds on each end was used. Unless otherwise indicated, all experiments were performed using *V. natriegens* crude cell extract incubated at 26°C for three hours, supplemented with 80 mM K-glutamate and 3.5 mM Mg-glutamate. The mean and standard deviations are shown (N=3).

Supporting Information

Attached supporting information document includes all materials & methods and supplementary figures for the main text.

Author contribution

DJW, NO, and HHL designed and performed experiments. DJW, NO, and HHL analyzed data and wrote the manuscript. GMC supervised the study.

Acknowledgments

The authors would like to thank Nina Donghia for helpful discussion regarding cell-free extract preparation and Dr. David Thompson for providing sfGFP standards.

Funding

This work was funded by the NIGMS 1U01GM110714-01 and DOE DE-FG02-02ER63445.

Notes

DJW, NO, and GMC have filed a patent related to this work.

References

- (1) Lee, H. H., Ostrov, N., Wong, B. G., Gold, M. A., Khalil, A., and Church, G. M. (2016, January 1) *Vibrio natriegens*, a new genomic powerhouse. *bioRxiv*.
- (2) Weinstock, M. T., Heseck, E. D., Wilson, C. M., and Gibson, D. G. (2016) *Vibrio natriegens* as a fast-growing host for molecular biology. *Nat. Methods*.
- (3) Fernández-Llamas, H., Castro, L., Blázquez, M. L., Díaz, E., and Carmona, M. (2017) Speeding up bioproduction of selenium nanoparticles by using *Vibrio natriegens* as microbial factory. *Sci. Rep.* 7, 16046.
- (4) Dalia, T. N., Hayes, C. A., Stolyar, S., Marx, C. J., McKinlay, J. B., and Dalia, A. B. (2017) Multiplex Genome Editing by Natural Transformation (MuGENT) for Synthetic Biology in *Vibrio natriegens*. *ACS Synth. Biol.*
- (5) Aiyar, S. E., Gaal, T., and Gourse, R. L. (2002) rRNA promoter activity in the fast-growing bacterium *Vibrio natriegens*. *J. Bacteriol.* 184, 1349–1358.
- (6) Hoffart, E., Grenz, S., Lange, J., Nitschel, R., Müller, F., Schwentner, A., Feith, A., Lenfers-Lücker, M., Takors, R., and Blombach, B. (2017) High substrate uptake rates empower *Vibrio natriegens* as production host for industrial biotechnology. *Appl. Environ. Microbiol.*
- (7) Long, C. P., Gonzalez, J. E., Cipolla, R. M., and Antoniewicz, M. R. (2017) Metabolism of the fast-growing bacterium *Vibrio natriegens* elucidated by (13)C metabolic flux analysis. *Metab. Eng.* 44, 191–197.
- (8) Li, J., Wang, H., Kwon, Y.-C., and Jewett, M. C. (2017) Establishing a high yielding streptomyces-based cell-free protein synthesis system. *Biotechnol. Bioeng.* 114, 1343–1353.
- (9) Wang, H., Li, J., and Jewett, M. C. (2018) Development of a *Pseudomonas putida* cell-free protein synthesis platform for rapid screening of gene regulatory elements. *Synth. Biol.* 3.
- (10) Moore, S. J., Lai, H.-E., Needham, H., Polizzi, K. M., and Freemont, P. S. (2017) *Streptomyces venezuelae* TX-TL - a next generation cell-free synthetic biology tool. *Biotechnol.*




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

J. 12.

- (11) Kim, J. K., Kang, H., Chae, J. S., Park, Y. H., and Choi, Y. J. (2000) Synthesis of cefminox by cell-free extracts of *Streptomyces clavuligerus*. *FEMS Microbiol. Lett.* *182*, 313–317.
- (12) Caschera, F., and Noireaux, V. (2014) Synthesis of 2.3 mg/ml of protein with an all *Escherichia coli* cell-free transcription-translation system. *Biochimie* *99*, 162–168.
- (13) Carlson, E. D., Gan, R., Hodgman, C. E., and Jewett, M. C. (2012) Cell-free protein synthesis: applications come of age. *Biotechnol. Adv.* *30*, 1185–1194.
- (14) Forster, A. C., and Church, G. M. (2007) Synthetic biology projects in vitro. *Genome Res.* *17*, 1–6.
- (15) Dudley, Q. M., Karim, A. S., and Jewett, M. C. (2015) Cell-free metabolic engineering: biomanufacturing beyond the cell. *Biotechnol. J.* *10*, 69–82.
- (16) PRATT, and M, J. (1984) Coupled transcription-translation in prokaryotic cell-free systems. *Transcription and translation: a practical approach.* 179–209.
- (17) Kwon, Y.-C., and Jewett, M. C. (2015) High-throughput preparation methods of crude extract for robust cell-free protein synthesis. *Sci. Rep.* *5*, 8663.
- (18) Sun, Z. Z., Hayes, C. A., Shin, J., Caschera, F., Murray, R. M., and Noireaux, V. (2013) Protocols for implementing an *Escherichia coli* based TX-TL cell-free expression system for synthetic biology. *J. Vis. Exp.* e50762.
- (19) Eagon, R. G. (1962) *Pseudomonas natriegens*, a marine bacterium with a generation time of less than 10 minutes. *J. Bacteriol.* *83*, 736–737.
- (20) Zawada, J., and Swartz, J. (2006) Effects of growth rate on cell extract performance in cell-free protein synthesis. *Biotechnol. Bioeng.* *94*, 618–624.
- (21) Sun, Z. Z., Hayes, C. A., Shin, J., Caschera, F., Murray, R. M., and Noireaux, V. (2013) Protocols for implementing an *Escherichia coli* based TX-TL cell-free expression system for synthetic biology. *J. Vis. Exp.* e50762.
- (22) Kwon, Y.-C., and Jewett, M. C. (2015) High-throughput preparation methods of crude extract for robust cell-free protein synthesis. *Sci. Rep.* *5*, 8663.
- (23) Failmezger, J., Rauter, M., Nitschel, R., Kraml, M., and Siemann-Herzberg, M. (2017) Cell-free protein synthesis from non-growing, stressed *Escherichia coli*. *Sci. Rep.* *7*, 16524.
- (24) Yun, H. S., Hong, J., and Lim, H. C. (1996) Regulation of ribosome synthesis in *Escherichia coli*: effects of temperature and dilution rate changes. *Biotechnol. Bioeng.* *52*, 615–624.
- (25) Iskakova, M. B., Szaflarski, W., Dreyfus, M., Remme, J., and Nierhaus, K. H. (2006) Troubleshooting coupled in vitro transcription-translation system derived from *Escherichia coli* cells: synthesis of high-yield fully active proteins. *Nucleic Acids Res.* *34*, e135.
- (26) Failmezger, J., Scholz, S., Blombach, B., and Siemann-Herzberg, M. (2018) Cell-Free Protein Synthesis From Fast-Growing *Vibrio natriegens*. *Front. Microbiol.* *9*, 1146.
- (27) Kelwick, R., Webb, A. J., MacDonald, J. T., and Freemont, P. S. (2016) Development of a *Bacillus subtilis* cell-free transcription-translation system for prototyping regulatory elements. *Metab. Eng.* *38*, 370–381.
- (28) Sun, Z. Z., Yeung, E., Hayes, C. A., Noireaux, V., and Murray, R. M. (2014) Linear DNA for rapid prototyping of synthetic biological circuits in an *Escherichia coli* based TX-TL cell-free system. *ACS Synth. Biol.* *3*, 387–397.
- (29) Seki, E., Matsuda, N., Yokoyama, S., and Kigawa, T. (2008) Cell-free protein synthesis system from *Escherichia coli* cells cultured at decreased temperatures improves productivity by decreasing DNA template degradation. *Anal. Biochem.* *377*, 156–161.
- (30) Michel-Reydellet, N., Calhoun, K., and Swartz, J. (2004) Amino acid stabilization for cell-

- 1
2
3 free protein synthesis by modification of the Escherichia coli genome. *Metab. Eng.* 6, 197–203.
4 (31) Wang, H., Li, J., and Jewett, M. C. (2018) Development of a Pseudomonas putida cell-free
5 protein synthesis platform for rapid screening of gene regulatory elements. *Synth. Biol.* 3.
6 (32) Li, J., Wang, H., and Kwon, Y. C. (2017) Establishing a high yielding streptomyces-based
7 cell-free protein synthesis system. *BioTechnology* .
8 (33) Sitaraman, K. (2004) A novel cell-free protein synthesis system. *J. Biotechnol.* 110, 257–
9 263.
10 (34) Shin, J., and Noireaux, V. (2010) Efficient cell-free expression with the endogenous E. Coli
11 RNA polymerase and sigma factor 70. *J. Biol. Eng.* 4, 8.
12 (35) Caschera, F., and Noireaux, V. (2014) Synthesis of 2.3 mg/ml of protein with an all
13 Escherichia coli cell-free transcription–translation system. *Biochimie* 99, 162–168.
14 (36) Gesteland, R. F. (1966) Isolation and characterization of ribonuclease I mutants of
15 Escherichia coli. *J. Mol. Biol.* 16, 67–84.
16 (37) Spirin, A. S., and Swartz, J. R. (2014) Cell-free Protein Synthesis: Methods and Protocols.
17 John Wiley & Sons.
18 (38) Chen, H., Xu, Z., Wang, C., Liao, Y., and Cen, P. (2008) Preliminary study on preparation
19 of E. coli cell-free system for protein expression. *Front. Chem. Eng. Chin.* 2, 224–229.
20 (39) Sitaraman, K. (2004) A novel cell-free protein synthesis system. *J. Biotechnol.* 110, 257–
21 263.
22 (40) Michel-Reydellet, N., Woodrow, K., and Swartz, J. (2005) Increasing PCR fragment
23 stability and protein yields in a cell-free system with genetically modified Escherichia coli
24 extracts. *J. Mol. Microbiol. Biotechnol.* 9, 26–34.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20



Vibrio natriegens

ACS Paragon Plus Environment
Cell extract

Cell free TX-TL